

Modelling functional responses with multivariate covariates

Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page



Page 1 of 32

Go Back

Full Screen

Close

Quit

1. Predicting temperature curves from climate zones

- We have 35 weather stations distributed across four climate zones:
 - Atlantic (16)
 - Pacific (6)
 - Continental (13)
 - Arctic (4)
- The dependent variable is $\text{Temp}(t)$, a function representing daily temperatures averaged over 1960–1994.
- The temperature functions were obtained by expanding the original 365 discrete daily averages in terms of 65 Fourier basis functions.

Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀ ▶

◀ ▶

Page 2 of 32


Go Back

Full Screen

Close

Quit

Montreal's temperature profile



Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀

▶

◀

▶

Page 3 of 32

Go Back

Full Screen

Close

Quit

The functional ANOVA model

- The model is

$$\text{Temp}_{mg}(t) = \mu(t) + \alpha_g(t) + \epsilon_{mg}(t).$$

- μ is the grand mean function
- α_g are the specific effects on temperature of being in climate zone g . To be able to identify them uniquely, we require that they satisfy the constraint

$$\sum_g \alpha_g(t) = 0 \text{ for all } t. \quad (1)$$

- ϵ_{mg} is the residual function showing unexplained variation specific to the k th weather station within climate group g .

Setting up the model

- Set up a 35 by 5 matrix \mathbf{Z} . Column 1 contains all 1's, and columns $g + 1, g = 1, \dots, 4$ contain zeros except for 1's in rows corresponding to stations in climate zone g .
- Append a final row with 0 in column 1, and 1's in the remaining columns.
- Let the functional response vector $\mathbf{Temp}(t)$ contain the 35 temperature profiles *plus* a final function that is zero for all t .
- Let functional regression coefficient vector $\beta(t)$ contain the functions $(\mu, \alpha_1, \dots, \alpha_4)$.
- The model in matrix notation,, including the zero sum constraint,is

$$\mathbf{Temp}(t) = \mathbf{Z}\beta(t) + \epsilon(t),$$

Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀ ▶

◀ ▶

Page 5 of 32

Go Back

Full Screen

Close

Quit

Fitting the model





- The residual $\text{Temp}_i(t) - \mathbf{Z}_i\boldsymbol{\beta}(t)$ is now a function.
- The least squares fitting criterion becomes

$$\text{LMSSE}(\boldsymbol{\beta}) = \sum_g^4 \sum_m^{N_g} \int [\text{Temp}_{mg}(t) - \sum_j^q z_{(mg),j} \beta_j(t)]^2 dt.$$

- This is minimized with respect to the regression functions by

$$\hat{\boldsymbol{\beta}}(t) = (\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{Z}'\mathbf{Temp}(t)$$

The region effects $\alpha_g(t)$



Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀ ▶

◀ ▶

Page 7 of 32

Go Back

Full Screen


Close

Quit


The mean plus region effects

$$\mu(t) + \alpha_g(t)$$


Atlantic




Pacific



Continental



Arctic



Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀

▶

◀

▶

Page 8 of 32


Go Back

Full Screen

Close

Quit

The standard error of measurement function $\sigma(t)$



Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀◀

▶▶

◀

▶

Page 9 of 32

Go Back

Full Screen

Close

Quit

2. Assessing fit

- Is there significant variation in temperature over climate zones? Of course there is! This does not seem like an interesting question.
- On the other hand, whether the Atlantic, Pacific and Continental stations are significantly different in the summer might be.
- Interesting summaries of fit, of effects, and inferences are likely to be *local* in nature.

[Home Page](#)[Title Page](#)[◀](#)[▶](#)[◀](#)[▶](#)

Page 10 of 32

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- It is useful to use the error sum of squares function

$$\text{SSE}(t) = \sum_{mg} [\text{Temp}_{mg}(t) - \mathbf{Z}_{mg}\hat{\boldsymbol{\beta}}(t)]^2.$$

to assess fit at or near time t .

- As in ordinary regression, we can compare this to the variation of the response about its mean

$$\text{SSY}(t) = \sum_{mg} [\text{Temp}_{mg}(t) - \hat{\mu}(t)]^2$$

- The corresponding mean squared error functions are

$$\text{MSE}(t) = \text{SSE}(t) / \text{df}(\text{error})$$

$$\text{MSR}(t) = \frac{\text{SSY}(t) - \text{SSE}(t)}{\text{df}(\text{model})}$$

Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀ ▶

◀ ▶

Page 11 of 32

Go Back

Full Screen

Close

Quit

Multiple correlation and F-ratio functions



- The squared multiple correlation function is

$$RSQ(t) = [SSY(t) - SSE(t)] / SSY(t).$$

- and the F-ratio function is

$$FRATIO(t) = \frac{MSR(t)}{MSE(t)}.$$

R^2 and F -ratio plots



Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀

▶

◀

▶

Page 13 of 32

Go Back

Full Screen

Close

Quit

3. Estimating the regression functions

$$\beta_j(t)$$

- We want a general framework for estimating functional parameters in this and other linear models.
- We want to be able to penalize the roughness of any parameter β_j .
- We also want the capacity to estimate confidence intervals for a parameter,
- and for functionals $\rho(\beta_j)$ of a parameter.

Home Page

Title Page

◀ ▶

◀ ▶

Page 14 of 32

Go Back

Full Screen

Close

Quit

Some basis function expansions for

$$\beta_j(t)$$

- Let the regression coefficient vector $\beta(t)$ have the expansion

$$\beta(t) = \mathbf{B}\theta(t)$$

where matrix \mathbf{B} is q by K_β and the K_β basis functions $\theta_\ell(t)$ are contained in vector $\theta(t)$.

- In the temperature example, it would be natural to use a certain number K_β of Fourier basis functions.

Home Page

Title Page

◀ ▶

◀ ▶

Page 15 of 32

Go Back

Full Screen

Close

Quit

A roughness penalty for $\beta_j(t)$

- If the response curves in $\mathbf{y}(t)$ are rough, we may want to impose some smoothness on the estimated β_j 's.
- Let L be a linear differential operator, such as $L = D^2$, that defines variation $L\beta(t)$ that we wish to penalize.
- Our roughness penalty on $\beta(t)$ is

$$\text{PEN}(\beta) = \int [L\beta(s)]' [L\beta(s)] ds .$$

Home Page

Title Page

◀ ▶

◀ ▶

Page 16 of 32

Go Back

Full Screen

Close

Quit

The penalized least squares criterion

- Let the response function vector $\mathbf{y}(t)$ have the basis function expansion in terms of K_y basis functions $\phi_k(t)$:

$$\mathbf{y}(t) = \mathbf{C}\phi(t)$$

- Then the penalized least squares function is

$$\begin{aligned} \text{PENSSE}(y|\beta) = & \int (\mathbf{C}\phi - \mathbf{ZB}\theta)' \mathbf{W} (\mathbf{C}\phi - \mathbf{ZB}\theta) \\ & + \lambda \int (\mathbf{LB}\theta)' (\mathbf{LB}\theta) . \end{aligned}$$

Penalized least squares in matrix terms

- we need to define these three matrices:

$$\mathbf{J}_{\phi\phi} = \int \phi\phi' , \quad \mathbf{J}_{\theta\theta} = \int \theta\theta' , \quad \mathbf{J}_{\phi\theta} = \int \phi\theta'$$

- and this roughness penalty matrix

$$\mathbf{R} = \int (L\theta)(L\theta)' .$$

- The fitting criterion now can be expressed as

$$\text{PENSSE}(y|\beta) = \text{trace}(\mathbf{C}'\mathbf{C}\mathbf{J}_{\phi\phi}) + \text{trace}(\mathbf{Z}'\mathbf{Z}\mathbf{B}\mathbf{J}_{\theta\theta}\mathbf{B}') - 2 \text{trace}(\mathbf{B}\mathbf{J}_{\theta\theta}\mathbf{C}'\mathbf{Z}) + \lambda \text{trace}(\mathbf{B}\mathbf{R}\mathbf{B}') ,$$

[Predicting temperature...](#)[Assessing fit](#)[Estimating the...](#)[Functional probes or...](#)[Correlations between...](#)[Summary](#)[Home Page](#)[Title Page](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 18 of 32](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

The normal equations for \mathbf{B}

- Taking the matrix derivative with respect to \mathbf{B} and setting it to 0 gives

$$(\mathbf{Z}'\mathbf{Z}\mathbf{B}\mathbf{J}_{\theta\theta} + \lambda\mathbf{B}\mathbf{R}) = \mathbf{Z}'\mathbf{C}\mathbf{J}_{\phi\theta}.$$

- We can use the Kronecker product to convert expressions of the form \mathbf{ABC}' to

$$\text{vec}(\mathbf{ABC}') = (\mathbf{C} \otimes \mathbf{A})\text{vec}(\mathbf{B}),$$

and consequently the normal equations become

$$[\mathbf{J}_{\theta\theta} \otimes (\mathbf{Z}'\mathbf{Z}) + \mathbf{R} \otimes \lambda\mathbf{I}]\text{vec}(\mathbf{B}) = \text{vec}(\mathbf{Z}'\mathbf{C}\mathbf{J}_{\phi\theta}).$$

- The estimate $\hat{\mathbf{B}}$ is therefore

$$\begin{aligned}\text{vec}(\hat{\mathbf{B}}) &= [\mathbf{J}_{\theta\theta} \otimes (\mathbf{Z}'\mathbf{Z}) + \mathbf{R} \otimes \lambda\mathbf{I}]^{-1}(\mathbf{J}_{\phi\theta} \otimes \mathbf{Z}')\text{vec}(\mathbf{C}) \\ &= \mathbf{S}_{\beta}\text{vec}(\mathbf{C}).\end{aligned}$$

[Predicting temperature...](#)[Assessing fit](#)[Estimating the...](#)[Functional probes or...](#)[Correlations between...](#)[Summary](#)[Home Page](#)[Title Page](#)[<<](#)[>>](#)[<](#)[>](#)[Page 19 of 32](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

4. Functional probes or contrasts

- Estimating the entire regression function $\beta_j(t)$ is fine,
- but we want to focus our attention on local or specific shape features of $\beta_j(t)$.
- Perhaps, for example, we want to examine the behavior of the temperature coefficient functions in mid-winter.
- A functional *probe* or *contrast* is of the form

$$\rho(\beta) = \int \xi(s)\beta_j(s) ds$$

- $\xi(s)$ is a weight function that we choose so as to concentrate our attention on a local region, or to look for specific patterns of variation in $\beta_j(t)$.
- There no particular need for $\xi(s)$ to integrate to 0.

- When $\beta_j(s)$ has the basis function expansion

$$\beta_j(s) = \mathbf{B}_j \boldsymbol{\theta}(s),$$

where \mathbf{B}_j is the j th row of \mathbf{B} , the contrast becomes

$$\rho(\beta) = \mathbf{B}_j \int \xi(s) \boldsymbol{\theta}(s) ds$$

Some examples

- *Point evaluation:*

$$\xi(s) = \delta(s - t)$$

This simply produces the function value $\beta(t)$.

- *Local behavior.* Assuming that β is periodic, we can use

$$\xi(s) = \exp[(s - t)^2 / (2\sigma)]$$

to assess the behavior of β in a neighborhood of t of a size determined by constant σ .

[Predicting temperature...](#)[Assessing fit](#)[Estimating the...](#)[Functional probes or...](#)[Correlations between...](#)[Summary](#)[Home Page](#)[Title Page](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 22 of 32](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

How do I work out confidence limits for these probes?

- The random element in a linear model is the residual function value

$$\epsilon_i(t_j) = y_{ij} - x_i(t_j).$$

- Any linear function of the data inherits it's variance from the variance of the data.
- The variance of the data conditional on the model is the variance of the residuals.

- We have two tasks:

- Estimate the variance of the residuals for a single response. (The mean can usually be taken to be 0.) Let's call this Σ_e .
- Assuming independence of the observations, the variance of the whole response data matrix is

$$\text{Var}[\text{vec}(\mathbf{Y})] = \Sigma_e \otimes \mathbf{I}.$$

- Work out the linear mapping from the data to the probe $\rho(\beta_j)$ that is being estimated. Let us call this \mathbf{M}_j .

- The rest is easy:

$$\text{Var}[\rho(\beta_j)] = \mathbf{M}_j'(\Sigma_e \otimes \mathbf{I})\mathbf{M}_j$$

Home Page

Title Page

◀ ▶

◀ ▶

Page 24 of 32

Go Back

Full Screen

Close

Quit

How do I work out mapping M_j ?

- In the examples given, $\rho(\beta_j)$ is three linear mappings removed from the data:

- The linear mapping from the raw data in matrix \mathbf{Y} to the coefficient matrix \mathbf{C} defining the smooth functions in $\mathbf{y}(t)$. This is

$$\text{vec}(\mathbf{C}) = (\mathbf{S}_y \otimes \mathbf{I})\text{vec}(\mathbf{Y}).$$

- The linear mapping from \mathbf{C} to the regression coefficient function coefficient vector \mathbf{B}'_j . We worked this out already, and called it \mathbf{S}_β .
- The linear mapping from \mathbf{B}'_j to the value of the probe. This is

$$\mathbf{U} = \int \xi(s) \boldsymbol{\theta}'(s) ds$$

Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀ ▶

◀ ▶

Page 25 of 32

Go Back

Full Screen

Close

Quit

- Now we have it, namely

$$\mathbf{M}_j = \mathbf{U}_j \mathbf{S}_\beta (\mathbf{S}_y \otimes \mathbf{I})$$

- This process is easy to extend to probes $\xi(s)$ involving all regression coefficients.
- For example, the variance of $\text{vec} [\hat{\beta}(\mathbf{t})]$ where \mathbf{t} is a vector of values of t , is

$$(\Theta \otimes \mathbf{I}) \mathbf{S}_\beta (\mathbf{S}_y \otimes \mathbf{I}) (\Sigma_e \otimes \mathbf{I}) (\mathbf{S}'_y \otimes \mathbf{I}) \mathbf{S}'_\beta (\Theta \otimes \mathbf{I})' .$$

where Θ is the matrix of values of θ at \mathbf{t} .

Some cautionary notes

- These sampling variances would only be “exact” if we knew Σ_e . The value of our confidence limit estimates depends critically on the quality of the estimate of Σ_e . There are many open questions about how to do this.
- We are assuming that the distribution of a probe is well summarized by its mean and variance.
- Our estimates are all conditioned on how many basis functions we use for both $y_i(t)$ and $\beta_j(t)$, namely K_y and K_β . Since we never know exactly how many to use, these should be regarded as random quantities, and a Bayesian treatment seems to be indicated.
- We should back up the use of these “delta method” confidence regions by bootstrapping and simulations.


Predicting temperature...
Assessing fit
Estimating the...
Functional probes or...
Correlations between...
Summary

[Home Page](#)[Title Page](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 27 of 32](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

5. Correlations between temperature and log precipitation residuals

- Once we have removed the climate zone effects from the temperature and log precipitation curves, are there one or more modes of correlation between them?
- We can use canonical correlation analysis to explore this question.
- We used the harmonic acceleration operator to impose smoothness on the two sets of canonical weighting functions.
- The smoothing parameter for the temperature residual was $\lambda = 10^7$, and for the log precipitation residual it was $\lambda = 10^9$.

The canonical correlations



Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀

▶

◀

▶

Page 29 of 32



Go Back

Full Screen

Close

Quit

The first two canonical weight functions



Predicting temperature...

Assessing fit

Estimating the...

Functional probes or...

Correlations between...

Summary

Home Page

Title Page

◀ ▶

◀ ▶

Page 30 of 32


Go Back

Full Screen

Close

Quit

The first two canonical variable scores



Predicting temperature ...

Assessing fit

Estimating the ...

Functional probes or ...

Correlations between ...

Summary

Home Page

Title Page

◀

▶

◀

▶

Page 31 of 32

Go Back

Full Screen

Close

Quit

6. Summary

- Regressing a functional response on multivariate independent variables or on a design matrix is not much different from the conventional regression analysis.
- One important difference is that we want to do *local* inference and interval estimation.
- We have, too, the capacity to smooth estimated functional parameters.
- But the number of basis functions that we use is not a fixed parameter in the traditional sense.

Home Page

Title Page

◀ ▶

◀ ▶

Page 32 of 32

Go Back

Full Screen

Close

Quit